

22-4-10

with TGR™ *Poa annua* Control



FEATURES & USE TIPS:

- Suppresses *Poa annua* growth in bentgrass, zoysiagrass, Kentucky bluegrass, perennial ryegrass and Kentucky bluegrass/perennial ryegrass fairways, tees and roughs
- Encourages preferential and aggressive growth of bentgrass, zoysiagrass, Kentucky bluegrass, and perennial ryegrass into adjacent *Poa annua* areas
- Extends color response through controlled-release nitrogen feeding and growth modification
- The growth and competitive ability of *Poa annua* will be reduced within 1-2 weeks of an application
- Sized (SGN 150) for easy spreading on fairways
- 50% of total nitrogen is from Poly-S slow release nitrogen

PRODUCT SPECIFICATIONS:

N (22%):	1.56% Ammoniacal Nitrogen 20.44% Urea Nitrogen *11.0% Slowly available nitrogen from polymer coated sulfur coated urea.
P - as P₂O₅ - (4%)	Ammonium Phosphate
K - as K₂O - (10%):	Potassium Sulfate
S - (1.02%):	Free Sulfur
Fe - (5%):	Iron
Active Ingredient:	Paclobutrazol (0.25%)

Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.

PRODUCT CODE:

AGC221PA5

Date: 011509
Supersedes: 042705
Label: L8

PRODUCT CHARACTERISTICS:

Product Type:

Combination Fertilizer with TGR™ *Poa annua* Control

Nitrogen Type:

Urea

Particle Size:

SGN: 150

Net Weight:

50 lbs.

Coverage:

10,800 sq. ft. @
Medium Rate

Andersons Golf Products and TGR are registered trademarks of The Andersons, Inc.

To order, contact your local Andersons Golf Products distributor.

For more information, contact Andersons at our website: www.andersonsgolfproducts.com or call 1-800-225-2639.

PRODUCT SHEET



22-4-10

with TGR® *Poa annua* Control

- Suppresses *Poa annua* growth in bentgrass, zoysiagrass, Kentucky bluegrass, perennial ryegrass and Kentucky bluegrass/perennial ryegrass fairways, tees and roughs
- Encourages preferential and aggressive growth of bentgrass, zoysiagrass, Kentucky bluegrass, and perennial ryegrass into adjacent *Poa annua* areas

ACTIVE INGREDIENT: 0.25%
OTHER INGREDIENTS: 99.75%
 Total 100.00%

TCASE# 6738-82-0

EPA Reg. No. 9199215
 EPA Est. Nos. 9199-0H-1M, 9199-0H-2E, 9199-AL-001A
 Underlined letter is first letter used in our code on bag.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Causes eye irritation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

	FRST AID
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If swallowed	Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance. Give gentle artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-752-8851 for emergency medical treatment information.	

Personal Protective Equipment
 Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof material, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions, for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

User Safety Recommendations
 Users should:
 • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
 • Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
 • Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply to steep slopes near water or when weather conditions favor drift from target areas. Do not contaminate water when disposing of equipment washwaters.

Guaranteed Analysis

Total Nitrogen (N)	22.00%
1.56% Ammoniacal Nitrogen	
20.44% Urea Nitrogen	
Available Phosphate (P ₂ O ₅)	4.00%
Soluble Phosphorus (K ₂ O)	10.00%
Sulfur (S)	1.02%
Iron (Fe)	5.00%

Plant nutrients derived from ammonium phosphate, urea, polymer coated sulfur coated urea, potassium sulfate and ferrous oxide.
 *11.00% slowly available nitrogen from polymer coated sulfur coated urea.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-ap.com>

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For all requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
 PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical resistant gloves made of any waterproof material such as barrier laminate or nitrile, nitrile or neoprene rubber or polyethylene and Stimp socks.

NON-AGRICULTURAL USE REQUIREMENTS
 This information is for non-agricultural uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR 170.10). The Worker Protection Standard applies when this product is used to produce agriculture plants on farms, forests, nurseries, or greenhouses.
 Do not allow people (other than applicator or pest on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

POA ANNUA CONTROL
 For suppression of *Poa annua* in bentgrass, zoysiagrass, Kentucky bluegrass, perennial ryegrass, and Kentucky bluegrass/perennial ryegrass fairways, tees and roughs. Repeat applications in a programmed approach will result in the gradual elimination of *Poa annua* as a major component of the grass stand. Use the desired rate of *Poa annua* control, annual applications and appropriate management practices will help prevent *Poa annua* from reestablishing as a major management problem.

What to expect
 The growth and competitive ability of *Poa annua* will be reduced within 1-2 weeks of an application. The stem and leaf tissue will become discolored (yellow to brown) for 3-8 weeks following the onset of growth regulation. Growth reduction of *Poa annua* will last for at least 3-5 weeks. During this period, growth of bentgrass, zoysiagrass, Kentucky bluegrass, and perennial ryegrass will be stimulated to "crowd out" the weakened *Poa annua*. Ongoing bentgrass will be more effective than Colonial bentgrass varieties in aggressively encroaching *Poa annua* areas. The formation of *Poa annua* seedheads will not be prevented, but spring and previous fall applications will stunt the growth of the seedhead stalk. Applications made just prior to seedhead emergence will greatly reduce the visibility of seedheads for 3-5 weeks. Regrowth of *Poa annua* will occur 4-8 weeks after application. Bentgrass, Kentucky bluegrass, and perennial ryegrass color will be enhanced for 6-12 weeks under non-stress conditions. Stress conditions following application may result in temporary undesirable color changes in bentgrass and Kentucky bluegrass. Avoid these conditions by irrigating and applying at the recommended time. Treatments will not have any detrimental effects on turfgrass root growth under normal growing conditions and when used as directed. The degree of *Poa annua* control will be inversely related to turfgrass management practices, climate, soil type, turf variety, and *Poa annua* biotype.

Program Scheduling
 Use any time when *Poa annua* is actively growing. Avoid applications during stressful conditions since the extent of discoloration on *Poa annua* and other desirable grasses may be unacceptable.

Fairways and roughs: Repeat late summer/early fall followed by spring applications are recommended for gradual *Poa annua* control. Avoid applications during stressful conditions.
Tees: Apply one or two weeks prior to *Poa annua* seedhead emergence to reduce seedhead visibility and encourage turf growth and coverage. Repeat applications can be made under non-stress conditions at least 10 weeks after *Poa annua* has recovered from any discoloration.

For long-term control in areas containing a high percentage of *Poa annua* contamination, repeat applications over a 2 to 3 year period may be required before the desirable grasses predominate. Moderate soil moisture conditions should be present before and after applications to achieve the best regulating effect (see precautions concerning saturated soil conditions). For best results, avoid applications during extreme soil temperature and moisture conditions. Spring applications should be delayed until any observable effects from pre-emergent control applications on bentgrass growth, color and quality have subsided. Where large areas of the desirable turf have been thinned from winter-damage, diseases or insects, without application until desired fill-in and coverage of the turf stand is achieved. When *Poa annua* populations have been reduced to the desired level, annual late summer/early fall applications are recommended where continuous long-term control is desired.

How to use
 Apply to dry foliage. For best results, water in (at least 1/4 inch) within 48 hours after application.

Recommended Application Rate
 LOW - for less overall discoloration of *Poa annua*
 MEDIUM - for quicker removal
 HIGH (50-70%) *Poa annua* population density
 Low (less than 50%) *Poa annua* population density

Additional Program Scheduling
 Overseeding is recommended to hasten conversion from *Poa annua* to the more desirable grasses. To avoid stunting growth of the desirable seedlings allow at least 2 weeks following treatment before overseeding, and in newly overseeded areas, make treatments at least 5 weeks after overseeding. If crabgrass or other annual weeds have been a problem in the past, an application of the appropriate pre-emergent control product should be made before using this product. If a weed, disease or insect problem occurs after application, apply a recommended control product in the same manner as is normally practiced, since the use of this product is compatible with the existing control products. If Embark® is used for

Poa annua seedhead control, apply this product at least 14 days after Embark® use to avoid potential discoloration of bentgrass and Kentucky bluegrass. Avoid late applications during onset of turfgrass dormancy to prevent fall and early spring discoloration of bentgrass and Kentucky bluegrass. Subsequent renewed growth and spring mowers will eliminate the potential discoloration. Applications made in the fall after the desirable grasses have ceased growth may result in less color response and reduced or delayed activity in affecting *Poa annua* but will provide both a greening response on the desirable grasses as well as a long-term suppression of *Poa annua* in the following spring. Calibrating clippings and the use of lightweight equipment throughout the year will enhance the long-term performance of treatments.

Embark® is a registered trademark of PBI Gordon Corporation.
Precautions
 Do not use on athletic fields where excessive wear is expected.
 Do not apply to Kentucky bluegrass collars and other areas around greens where consistent turf height and color are desired.

Not for use around shrubs, fruit trees, flowers or vegetable plants; however, applications to turf areas under the tree canopies will not affect or harm trees.
 Do not use during periods of extreme environmental stress, such as heat, drought, or cold, or during heavy insect or disease activity.

Frequent irrigation after application throughout the hot and dry weather conditions will help prevent potential discoloration of bentgrass and Kentucky bluegrass and ensure continued growth of the desirable grasses in *Poa annua* areas.

Heavy rainfall or irrigation after application in areas where the soil is already saturated may cause the active ingredient to move laterally on steep slopes and collect in low areas. These areas may undergo more severe growth control for a longer period of time. To avoid this response, do not apply when soil is already saturated.

Do not use spreader settings other than those recommended. Inproper rates may cause undesirable turf growth control and areas may discolor temporarily.
 When large areas of the desirable turf have been thinned from winter damage, disease, or insects, without application until desired fill-in and root-coverage of the turf stand is achieved.

Use in areas containing greater than 70% *Poa annua* will result in more widespread discoloration and may not appear acceptable to the user. Use of other cultural techniques are encouraged to lower the percentage of *Poa annua* below the 70% level.

GROWTH REGULATION OF LAWN TURFGRASSES
 For regulation of well-maintained, lawn-maintained, lawn-type commercial, industrial and residential bluegrass and ryegrass turfgrass areas that are on regular fertilizer programs.

What to Expect
 Applications made at the MEDIUM rate between March 1 and April 30 (Spring) or September 1 and October 31 (Fall) will result in reduced vertical growth, mowing frequencies and clipping yields of lawn-height bluegrass and ryegrass. *Poa annua* will be suppressed in treated turfgrass areas. Also treated will be improved and extended color and the overall improvement in turfgrass quality.

Precautions
 Do not make more than 2 spring and 1 fall applications.
 Do not make applications to turf areas containing more than 50% *Poa annua*.
 Do not use during the recommended period if extreme environmental stresses such as heat, drought or cold or if heavy insect or disease pressures are present.

STORAGE AND DISPOSAL
 Do not contaminate water, food, or feed by storage and disposal. **STORAGE:** Store in a clean, dry place. Re-seal opened bag by folding top down and securing. **PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary manner or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT - READ BEFORE USE:
 Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If items are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.


WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:
 Manufacturer makes no representation or description on the label and is not responsible for the results obtained from the use of this product in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS: WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS. YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION. (AP10044)

Product of USA
 Manufactured By:
 The Andersons Lawn Fertilizer Division, Inc.
 P.O. Box 114387
 Moline, IL 61702
www.andersonsfertilizer.com

Andersons Golf Products and TGR are registered trademarks of The Andersons, Inc.

SPREADER SETTINGS	MEDIUM RATE			LOW RATE			GROUND SPEED
	HALF WIDTH OF COVERAGE	SETTING	FULL WIDTH OF COVERAGE	SETTING	FULL WIDTH OF COVERAGE	SETTING	
Andersons AccuPro 2000, SR-2000 (one 8)	5.5 feet	M 1/2	11 feet	I	11 feet	K 3/4	3 mph
Scotts Rollax R-8, R-8A (cone 8)	5.5 feet	K 3/4	11 feet	H 3/4	11 feet	J 1/2	3 mph
Ley Models WTR, WFR, HR, 1250 (PTO at 450 rpm) Patrim III-B	19 feet	6 III-B	38 feet	3 1/2 III-B	38 feet	5 III-B	4.5 mph
Vicom (C3 Series)	15 feet	30	30 feet	18	30 feet	25	4.5 mph
APPLICATION INFORMATION							
Lb. Paclobutrazol/A							
Lb. product/1,000 sq. ft.							
Bogstare							
0.38							
3.5							
3.0							



0 31865 31128 4

SGN 150
Net Weight 50 lbs. (22.68 kg)
AGC221PA5
 L8 (AGCBLKPPUR29B)