

# Fertilizer with 1.07% ProPendi®



## 7-0-11

### FEATURES & USE TIPS:

- Provides preemergent control of crabgrass, goosegrass, foxtail, *Poa annua*, chickweed, oxalis, spurge, and other annual grassy and broadleaf weeds as listed.
- Provides a feeding of nitrogen and potassium.
- Provides initial greening plus controlled-release nitrogen.
- Includes 3.0 pound/acre active ingredient for certain hard to control weeds.

### PRODUCT SPECIFICATIONS:

N (7%):	Urea Nitrogen
K - as K <sub>2</sub> O - (11%):	Potassium Chloride
Cl (9.20%):	Chlorine Max
<b>Active Ingredient:</b>	Pendimethalin (1.07%)

*Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.*

### PRODUCT CODE:

## AGC72PD5

Date: 073108  
Supersedes: 000000  
Label: G8

### PRODUCT CHARACTERISTICS:

#### **Product Type:**

Combination fertilizer and Pendimethalin pre-emergent herbicide

#### **Nitrogen Type:**

NS-54™

#### **Particle Size:**

SGN: 215

#### **Net Weight:**

50 lbs.

#### **Coverage:**

13,100 sq. ft. @  
Low Rate

Andersons Golf Products and ProPendi are registered trademarks of The Andersons, Inc.; NS-54 is a trademark of The Andersons, Inc.

*To order, contact your local Andersons Golf Products distributor.*

*For more information, contact Andersons at our website: [www.andersonsgolfproducts.com](http://www.andersonsgolfproducts.com) or call 1-800-225-2639.*

# Andersons Golf Products® Fertilizer with 1.07% ProPendi®

## 7-0-11

### For Occupational Use on Golf Courses Only

- Provides preemergent control of crabgrass, goosegrass, foxtail, *Poa annua*, chickweed, oxalis, spurge and other annual grassy and broadleaf weeds.
- Provides a feeding of nitrogen and potassium
- Provides initial greening plus controlled-release nitrogen

## KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals:** CAUTION, causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Avoid inhalation. Do not allow people or pets to enter the treated area until dusts have settled. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed droppings to livestock.

#### Fish Aid

**Use:** Fish Aid is a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless instructed by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **Eye:** Rinse eyes with copious amounts of water for at least 15-20 minutes. Call a poison control center or doctor for treatment advice. **First Aid:** If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

#### Personal Protective Equipment

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protection equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash personal protection equipment separately from other laundry.

#### User Safety Recommendations:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE before leaving this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Engineering Controls

When handlers use closed systems, enclosed cabs, or encuffs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240 (d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

**Environmental Hazards:** This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, either collect for use or dispose of properly.

#### ACTIVE INGREDIENT:

Permethrin 1.07%

Inert Ingredients: 98.93%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agriculture workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box do not apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides, 40 CFR part 170. The Worker Protection Standard applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts have settled.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, tall fescue, fine fescue, bermuda grass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, centipede grass, zoysiagrass, bahiagrass, and zoysiagrass. See Limitation of Liability statements.

Not for use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bermudagrass-containing turf and *Poa annua* are the desired species, a light rate application (1.5 lbs. a/A) only is recommended.

#### How/When to Use

For control of annual grass and broadleaf weeds such as crabgrass, goosegrass, foxtail, lamyragrass, fall panicum, *Poa annua*, oxalis, knotweed, purslane, and prostrate spurge, apply in spring before they sprout.

To prevent *Poa annua* and fall germinating chickweed, apply product to dry foliage in late summer-early fall (before germination) at the RECOMMENDED RATE. Repeat application yearly for continued control. A repeat application made after *Poa annua* treatment should not be sooner than 4 months.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application (1/4-1/2 inch) to move granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not apply the chemical barrier between the soil surface by going or driving on a segment to application. Allow at least 4+ month span between the applications when repeat applications are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried. This product is not intended for use on dichondra or cool season turfgrasses where it is grown and maintained as the permanent stand.

When preemergence herbicide is applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be extensive in low-cut bermudagrass and on sandy soils.

Do not apply to dormant bentgrass or *Poa annua* (turf that is not actively growing or has experienced frost). As with most other preemergence herbicides, application of this product to bermudagrass overseeded with cool season turfgrasses species may require and thin overseeding.

On cool season turfgrasses and applications when temperatures are regularly in the 80's. This product may cause discoloration and thinning where over applications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature, below 40°F, seeping, and excessive soil moisture).

Do not apply to bentgrass and *Poa annua* after late summer.

Apply application to cement, stone, cobble or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (pail, course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

#### Properties of All Dintriazoline (DMA) Products

May cause temporary slowing of turf-in and stolon-rhizome rooting where turf has been thinned. Requires a waiting period before seeding or sprigging.

#### Seeding or Sprigging Intervals after Treatment

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

#### Treatment Interval after Seeding or Sprigging

On newly planted areas, reseedative planted turfgrasses has been mowed at least 4 times, and has filled in before making an application. On newly sprigged bermudagrass or other vegetative planted turfgrasses wait until the new grass has filled in and the stools have tilled down before application.

#### Notes

Do not apply more than a cumulative total of 280 lbs. of product (a 0.28 lbs. active ingredient) per year on bentgrass and *Poa annua*; 560 lbs. of product (a 0.56 lbs. active ingredient) per year on cool season turfgrasses; and 840 lbs. of product (a 0.84 lbs. active ingredient) per year on warm season turfgrasses.

TURFGRASS SPECIES	WEEDS	RATES OF THIS PRODUCT	COMMENTS
<b>Warm Season Grasses</b> Bermudagrass St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass (85% bs:acre rate or less) Tall Fescue  (Not for use on greens and tees)	Crabgrass Foxtail <i>Poa annua</i> Barnyardgrass Fall panicum Oxalis Prostrate spurge Purslane Knotweed Chickweed	163-280 lbs./acre prior to weed germination.  163-280 lbs./acre prior to weed germination.	Apply a repeat application of 140-163 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	Goosegrass	163-280 lbs./acre prior to weed germination.	Apply a repeat application of 140-163 lbs./acre in 6-8 weeks to control goosegrass if the other rate was used initially.
	Cudweed Hop clover Eveningprimrose Herbit	280 lbs./acre prior to weed germination.	

140 lbs. product/acre of 0.25 lbs./1,000 sq. ft. provides 1.50 lbs. active ingredient/acre and 0.22 lbs. nitrogen/1,000 sq. ft.  
163 lbs. product/acre of 0.25 lbs./1,000 sq. ft. provides 1.75 lbs. active ingredient/acre and 0.26 lbs. nitrogen/1,000 sq. ft.  
187 lbs. product/acre of 0.25 lbs./1,000 sq. ft. provides 2.00 lbs. active ingredient/acre and 0.30 lbs. nitrogen/1,000 sq. ft.  
280 lbs. product/acre of 0.30 lbs./1,000 sq. ft. provides 3.00 lbs. active ingredient/acre and 0.46 lbs. nitrogen/1,000 sq. ft.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

#### WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the technical description on the label and is, reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

**LEGAL RIGHTS:** THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/ JURISDICTION TO STATE/ JURISDICTION. (R16md103107)

Product of U.S.A.

Manufactured by:  
The Andersons Lawn Fertilizer Division, Inc.  
P.O. Box 119, Maumee, OH 43537  
www.andersonslawnproducts.com

Andersons Golf Products and ProPendi are registered trademarks of The Andersons, Inc.

#### Suggested Spreader Settings

To provide proper distribution, calibrate spreader before application.

50 lbs. treats 13,300 sq. ft. (0.35 acre) at the 140 lbs. product/acre - LIGHT RATE (3.25 lbs/1,000 sq. ft.)  
50 lbs. treats 15,100 sq. ft. (0.30 acre) at the 163 lbs. product/acre - LOW RATE (3.79 lbs/1,000 sq. ft.)  
50 lbs. treats 17,500 sq. ft. (0.26 acre) at the 187 lbs. product/acre - MEDIUM RATE (4.33 lbs/1,000 sq. ft.)  
50 lbs. treats 7,600 sq. ft. (0.17 acre) at the 280 lbs. product/acre - HIGH RATE (6.50 lbs/1,000 sq. ft.)

SREADER	GROUND SPEED	WIDTH OF COVERAGE	LIGHT RATE (0.30 lbs./acre)	LOW RATE (1.75 lbs./acre)	MEDIUM RATE (2.20 lbs./acre)	HIGH RATE (3.00 lbs./acre)
Andersons AccuPro 2000, SP2000, (cone 8)	3 mph	12 feet	L 1/4	L 3/4	M 1/4	N 3/4
Scotts R-8A (cone 8)	3 mph	12 feet	J	J 1/2	K	M
Lely Modus WTR/WR, HR, 1250 (Pail at 460 rpm), Pallet 11/8	4.5 mph	38 feet	5 1/4	5 3/4	6 1/4	7 1/4
Vicon (103 Series)	4.5 mph	32 feet	23	25	27	34

SGN 215

Net Wt 50 Lbs (22.63 kg)



# AGC72PD5

GB AGCBLKPDNA(B)

**DIRECTIONS FOR USE**  
It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Keep people and pets out of the area during application.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds	Dicot Weeds
Crabgrass	Cudweed
Foxtail	Hop clover
Barnyardgrass	Knweed
Goosegrass	Herbit
	Prostrate spurge
	Purslane
	Eveningprimrose

#### GUARANTEED ANALYSIS

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Total Potassium (K<sub>2</sub>O) 0.00%

Available Potassium (K<sub>2</sub>O) 0.00%

Total Nitrogen (N) 1.07%

Available Nitrogen (N) 0.983%

Total Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%

Available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.00%