


# Fungicide V



## FEATURES & USE TIPS:

- Prevents and controls *Pythium* blight on golf course greens and fairways (see label recommendations).
- Prevents and controls gray snow mold on golf course greens and fairways (see label recommendations).
- Convenient, dry-applied, ready-to-use, free-flowing product with active ingredient impregnated on and into a granular cob carrier.
- Choice of application rates depending upon the type and severity of infestation and degree of protection desired.
- May be applied by drop or rotary spreader.

## PRODUCT CODE:

**AGC87135.1**

Date: 061107  
Supersedes: 000000  
Label: G5

## PRODUCT CHARACTERISTICS:

### **Product Type:**

Fungicide

### **Net Weight:**

32.75 Lbs.

### **Coverage:**

11,000 Sq. Ft. @  
recommended rate

## PRODUCT SPECIFICATIONS:

**Active Ingredient:** Chloroneb (6.25%)

Andersons Golf Products is a trademark of The Andersons, Inc.

*To order, contact your local  
Andersons Golf Products  
distributor.*

*For more information, contact  
Andersons at our website:  
[www.andersonsgolfproducts.com](http://www.andersonsgolfproducts.com)  
or call 1-800-225-2639*

# Andersons Golf Products Fungicide V

- Prevents and controls Pythium blight & gray snow mold on golf course greens and fairways

## SGN 100

Covers up to 11,000 Sq. Ft.

Net Wt. 32.75 Lbs. (14.86 kg)

### ACTIVE INGREDIENT:

Chloroneb (1,4-dichloro-2,5-dimethoxybenzene) . . . 6.25%

INERT INGREDIENTS: . . . . . 93.75%

TOTAL . . . . . 100.00%

EPA Reg. No. 9198-182 EPA Est. No. 9198-OH-1M,  
9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code on bag.

### KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### PRECAUTIONARY STATEMENTS

**Hazards to Humans and Domestic Animals:** May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not graze or feed clippings from treated areas to livestock.

#### FIRST AID

<b>IF IN EYES</b>	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
<b>IF SWALLOWED</b>	• Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
<b>IF ON SKIN</b>	• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
<b>IF INHALED</b>	• Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

DOUBLE RATE prior to the first snow and at the RECOMMENDED RATE when snow melts in the Spring.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Call your local or state solid waste agency or call 1-800-CLEANUP if you have questions.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

### WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product con-

forms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

### LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/ JURISDICTION TO STATE/JURISDICTION. (RVamd061202)

Product of USA

Manufactured by:

The Andersons Lawn Fertilizer Division Inc.  
P. O. Box 119  
Maumee, OH 43537

Visit us at our website:

[www.AndersonsGolfProducts.com](http://www.AndersonsGolfProducts.com)

**AGC87135.1-G5** (AGCBLKPBLU.1B)

©2004

SUGGESTED SPREADER SETTINGS			
<i>To provide proper distribution calibrate your spreader before application.</i>			
32.75 lbs. treats 11,000 sq. ft. (0.25 acre) at the RECOMMENDED RATE (2.98 lbs./1,000 sq. ft.)			
32.75 lbs. treats 5,500 sq. ft. (0.13 acre) at the DOUBLE RATE (5.95 lbs./1,000 sq. ft.)			
SPREADER	GROUND SPEED	SPREADER SETTINGS	
		WIDTH OF COVERAGE	DOUBLE RATE
The Andersons AccuPro 1000	3.0 mph	10	0 1/2
AccuProSR-2000 (cone 9)	3.0 mph	10	0 1/2
Andersons Drop SS-2	3.0 mph	overlap wheels	8
Scotts Rotaries R-8, R-8A (cone 9)	3.0 mph	10	0 1/2
Lely W Models WTR, WFR, HR, 1250 (PTO at 450 rpm)	4.5 mph	16	6 III
Vicon (03 Series)	4.5 mph	16	20

#### SUGGESTED SPREADER SETTINGS

To provide proper distribution calibrate your spreader before application.

32.75 lbs. treats 11,000 sq. ft. (0.25 acre) at the RECOMMENDED RATE (2.98 lbs./1,000 sq. ft.)

32.75 lbs. treats 5,500 sq. ft. (0.13 acre) at the DOUBLE RATE (5.95 lbs./1,000 sq. ft.)

SPREADER	GROUND SPEED	SPREADER SETTINGS	
		WIDTH OF COVERAGE	DOUBLE RATE
The Andersons AccuPro 1000	3.0 mph	10	0 1/2
AccuProSR-2000 (cone 9)	3.0 mph	10	0 1/2
Andersons Drop SS-2	3.0 mph	overlap wheels	8
Scotts Rotaries R-8, R-8A (cone 9)	3.0 mph	10	0 1/2
Lely W Models WTR, WFR, HR, 1250 (PTO at 450 rpm)	4.5 mph	16	6 III
Vicon (03 Series)	4.5 mph	16	20

**Environmental Hazards:** Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

### Prevents and controls:

Pythium blight, gray snow mold. Whenever used on greens and fairways, apply to moist foliage in the morning or evening. Sprinkle lightly just before traffic resumes to avoid pickup of material on shoes, golf balls, or maintenance equipment.

**To prevent Pythium blight,** apply at the

RECOMMENDED RATE to moist foliage when high temperature and humidity favor development. Repeat at 5-7 day intervals as long as these weather conditions persist.

**To control Pythium blight,** apply at the

RECOMMENDED RATE to moist foliage as soon as disease activity is observed; repeat at 5-7 day intervals during high temperature and humidity conditions.

In case of severe disease attack, apply at the DOUBLE RATE to moist foliage and continue RECOMMENDED RATE applications at 5-7 day intervals during high temperature and humidity conditions.

On new seedlings, apply at the RECOMMENDED RATE after sowing seed; at first signs of disease, apply to moist foliage of newly germinated grass and repeat at 7-10 day intervals if evidence of disease persists. When treating to control Pythium blight, do not fertilize or water heavily until disease activity is arrested; prolonged contact with foliage enhances effectiveness.

**Gray snow mold:** Apply at the RECOMMENDED RATE to dry or moist foliage prior to the first heavy snow (early Winter). Repeat in mid-Winter if turf is free of snow. A final treatment in late Winter as the snow melts extends protection. In areas where snow cover is present the entire Winter, apply at the