

Turf Fertilizer 0-0-22

with Dimension® Turf Herbicide & Ronstar®



FEATURES & USE TIPS:

- Provides preemergent control or suppression of the following annual weeds: goosegrass, crabgrass, annual bluegrass, and foxtail.
- Labeled for use on established or sprigged bermudagrass, zoysiagrass and St. Augustinegrass.
- Utilizes Ronstar and Dimension herbicides to maximize performance benefits of each active ingredient.
- Uniformly sized granules minimize particle segregation, which contributes to more accurate distribution.
- The readily available potassium increases resistance to heavy traffic and other turf stresses.
- Small particle formula provides more particles per square inch for better herbicide distribution.

PRODUCT SPECIFICATIONS:

K - as K₂O (22%):	Potassium Sulfate
S (7.48%) :	Combined Sulfur
Active Ingredients:	0.125% Dithiopyr 1.000% Oxadiazon

PRODUCT CODE:

AGCDMR5

Date: 111307
Supersedes: 101702
Label: J7

PRODUCT CHARACTERISTICS:

Product Type:

Combination fertilizer,
Dimension® &
Ronstar® Herbicides

Particle Size:

SGN: 125

Net Weight:

50 lbs.

Coverage:

10,800 sq. ft. @
High rate

Dimension is a registered trademark of Dow AgroSciences L.L.C.; Ronstar is a registered trademark of Bayer; Andersons Golf Products is a trademark of The Andersons, Inc.

*To order, contact your local
Andersons Golf Products
distributor.*

*For more information, contact
Andersons at our website:
www.andersonsgolfproducts.com
or call 1-800-225-2639*



This product is a selective herbicide that provides control of listed annual grasses and broadleaf weeds in established lawns and ornamental turfs.

For sale to and use by professional applicators only. Not for sale to or use by homeowners/consumers.

Not for Use on Home Lawns
 New York Use Restrictions: In New York State this product may only be used by professional applicators and at no more than 400 pounds per acre (equal to 0.5 pound Dimension active ingredient per acre), and is prohibited from use in Nassau and Suffolk Counties.

GUARANTEED ANALYSIS

Soluble Potash (K ₂ O)	22.00%
Sulfur (S)	7.46%
7-46% Combined Sulfur	
Plant nutrients derived from potassium sulfate.	
F292	
ACTIVE INGREDIENTS:	
Oxadiazon	1.000%
Dithiopyr	0.125%
INERT INGREDIENTS:	98.875%
Total	100.000%

EPA Reg. No. 9198-154
 EPA Est. No. 9198-OH-IL 9198-OH-2B, 9198-AL-001A
 (Underlined letter is first letter used in run code on end of bag)

**KEEP OUT OF REACH OF CHILDREN
 CAUTION**

Some components of this product may cause eye irritation
PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

- Long-sleeved shirt and long pants, and
- Chemical resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for wash-ables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly, and change into clean clothing.

FIRST AID

- | | |
|--------------------------------|--|
| If on skin or clothing: | <ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice |
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contacts, if present, after the first 5 minutes; then continue rinsing • Call a poison control center or doctor for treatment advice |

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-3387 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. Use with care when applying to turf areas adjacent to any body of water. Drift and runoff

from treated turf may adversely affect organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from treated areas. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the entire label before using this product. Use only according to label instructions. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For most economical use, rates of this product should be based on anticipated weed population, time of application and length of growing season. A well-maintained turf with low weed infestations will require less herbicide for effective control than will a thin turf with low fertility, minimum maintenance and heavy traffic. Geographic areas with longer growing seasons require higher rates. When cultural practices require application of this product well in advance of weed seed germination, rates should be adjusted upward to compensate for normal breakdown of the active ingredient.

GENERAL INFORMATION

This product is a selective, preemergent herbicide for control of many annual grasses and broadleaf weeds in established lawns and ornamental turfs, especially golf courses, lawns and roughs. This product controls weeds by killing the young turf seedlings as they come in contact with the herbicide during germination, by a combination of two different mechanisms. The two herbicides affect different parts of the weeds, which aids in providing excellent weed control while minimizing adverse effects to the turf. This product will not harm newly established ornamentals when used according to label directions (avoid foliage contact).

USE PRECAUTIONS

Do not apply to wet turf as discoloration may occur. Do not allow domestic animals to graze treated areas. Avoid contamination of water intended for irrigation and domestic use.

TURFGRASS SAFETY: This product may be used on lawns and ornamental turfs that are well-established. The grass must have developed a good root system and a uniform stand, and have received at least two mowings following its seeding, sodding, or sprigging before it can receive its first application of this product. Use of this product on lawns and ornamental turfs that are not well-established or on those that have been weakened by weather, pest, disease, chemical, or mechanical-related stress, may increase the chances for turf injury.

This product should only be applied to lawns and ornamental turfs that are composed of the following turfgrass species that have been determined to be tolerant to applications of this product. When applied as directed under the conditions described, the following established turfgrasses are tolerant to this product:

- Warm-Season Grasses**
- Bermudagrass (*Cynodon dactylon*)
 - Centipedegrass (*Axonopus affinis*)
 - Kentuckygrass (*Pennisetum clandestinum*)
 - St. Augustinegrass (*Stenotaphrum secundatum*)
 - Zoysagrass (*Zoysa japonica*)

Red (Fine) Rescue and Bentgrass are not tolerant of this product. The use of this product on turf containing Red Fescue or Bentgrass should be limited to situations where they are undesirable grasses.

Do not apply to Dichondra, Centipedegrass or newly seeded areas.

RESEEDING, OVERSEEDING, OR SPRIGGING: Reseeding, overseeding, or sprigging of treated areas within 4 months after application of this product may inhibit the establishment of desirable turfgrasses. When reseeding or overseeding, proper cultural practices such as soil cultivation, irrigation and fertilization should be followed. For best results, use mechanical or power seeding equipment (sift seeders) designed to give good seed to soil contact.

GOLF COURSE PUTTING GREENS AND TEES: DO NOT use this product on golf course putting greens and tees because of the varying cultural practices used.

OTHER USE PRECAUTIONS: The early postemergence crabgrass control described below is limited. Postemergence applications are only effective on crabgrass if applied before crabgrass has displayed its fifth leaf or first tiller.

For best results, cultural practices that disturb the soil, such as core, spike, or hydro-aerification, and verticulating, should be done before application of this product. DO NOT apply this product until the grass has recovered from these cultural practices.

Apply this product directly to established lawns or ornamental turfs only. DO NOT allow product or drift to contact foliage of flowers, vegetables, shrubs, or trees.

DO NOT use clippings from treated turf for mulching around vegetables or fruit trees. DO NOT apply this product to pastures. Keep people and pets off treated areas until product has been thoroughly watered in and grass has dried.

APPLICATION DIRECTIONS

PREEMERGENCE CONTROL OF CROWFOOTGRASS (GOOSEGRASS) AND CRABGRASS: This product provides "preemergence" control of crowfootgrass (goosegrass) and crabgrass when applied prior to the emergence of these weeds from the ground in established lawns and ornamental turfs. Crowfootgrass germinates somewhat later than crabgrass.

This product can provide "early postemergence" suppression of crabgrass during the early stages of crabgrass growth after the crabgrass has emerged from the ground. However, it is often difficult to see the very small, early stages of crabgrass in well-established lawns and ornamental

turfs. Postemergence crabgrass suppression will be obtained only when this product is applied prior to the tillering of crabgrass, which generally corresponds to the time when you can first easily see the crabgrass plants in the lawn or turf. So to the practical benefit of this product's additional, early postemergence activity, it can be applied to newly emerging crowfootgrass and crabgrass products. It can give the user a 2-4 week longer period of time to make applications and still suppress crabgrass. This product is not intended to control large areas of established crabgrass, and must be used prior to crowfootgrass (goosegrass) and crabgrass control.

Application Frequency and Timing: For season-long control, this product should be applied as a single application in the spring for crowfootgrass (goosegrass) and crabgrass control.

DO NOT apply more than 400 lbs. per acre. (0.5 lbs. Dimension and 4.0 lbs. Ronstar Alacize) per application, and no more than 800 lbs. per acre (0.75 lbs. Dimension and 6.0 lbs. Ronstar Alacize) per year, except in areas where there is a heavy weed infestation. In areas of heavy weed infestation, do not apply more than 800 lbs. of product per acre (1.0 lbs. Dimension and 6.0 lbs. Ronstar Alacize) per year.

Spring Applications: For spring applications made in the spring, this product should be applied at the appropriate rate corresponding to one of the two control programs listed in Table 1 below, depending on the user's location, the turfgrass mowing height, and whether the use is considered to be preemergence or early postemergence at the time of the application. The duration of residual weed control provided by this product is directly related to the mow rate applied, but will vary somewhat depending on weather, weed pressure, turfgrass competitiveness, and the user's location within a region.

Use the Low Rate Program for preemergence control at sites where the turfgrass is cut high (e.g., golf courses), and so is especially useful in the transition zone. This program may be used in areas of light weed pressure, and is not for early postemergence crabgrass control.

Use the High Rate Program for preemergence control at sites where (a) turfgrass is cut low (e.g., golf fairways) and/or (b) turf maintenance or weed control has not been conducted during the previous year. This program provides 4-6 months of preemergence crowfootgrass (goosegrass) and crabgrass control, and so is especially useful in the southern and coastal south turf zones. **South of Orlando, or in areas that do not receive at least one frost which naturally controls crowfootgrass (goosegrass) and crabgrass, may require the use of a postemergence crabgrass product to eliminate existing weeds.** This product also provides early postemergence crabgrass suppression at sites where turfgrass is cut low (e.g., golf fairways).

TABLE 1: Recommended Single Application Use Rates (lbs/acre)

Low Rate	High Rate
150 lbs. Ronstar, 1.5 lbs. Al per acre	200 lbs. Ronstar, 2.0 lbs. Al per acre
Dimension: 0.1875 lbs. Al per acre	Dimension: 0.25 lbs. Al per acre

Area Classifications: DE, KS, KY, MD, MO, NJ, VA, southeastern PA, southern areas of IL, TN, OH, & coastal areas of CT, NY, & RI. **South:** AL, AR, AZ, CA, GA, LA, MS, NC, NM, NV, OK, SC, TN, & TX. **Coastal South:** HI, FL, & southern coastal areas of AL, GA, LA, MS, NC, SC, & TX.

This product may cause temporary discoloration of Bermudagrass and St. Augustinegrass. This is normally outgrown within 2-3 weeks.

Tips for Improved Control: For best results, apply this product within a few days after mowing and delay mowing again for a few days after the application. When treated lawn or ornamental turf areas are watered or receive significant rainfall within a few days after application of this product, improved weed control may result.

CONTROL OF OTHER GRASS AND BROADLEAF WEEDS

Spring Applications: When used as directed for crowfootgrass (goosegrass) and crabgrass control in the spring, this product will also control suppress the following weeds when applied prior to their emergence:

- annual bluegrass
- annual and perennial ryegrass
- barnyardgrass
- bitertgrass
- Carolina geranium
- carpetgrass
- chickweed
- common groundsel
- common knotweed
- common purslane
- corn speedwell
- foxtail (yellow & green)*
- Florida pursley
- herb
- ovals (buttercup, creeping & yellow woodsoorrel)
- parsley-piert
- pigweed
- pineappleweed
- smartweed
- spruce (prostrate & spotted)
- stinging nettle

APPLICATION EQUIPMENT AND INSTRUCTIONS

For best results, water-in the product as soon as practical after application. Thorough watering assures more even distribution of the herbicide and avoids gaps in coverage.

Apply this product with drop or rotary-type spreaders designed to apply granular herbicides. Avoid the use of spreaders that would apply this product in narrow rows or concentrated bands. Before each application, calibrate the spreader according to the equipment manufacturer's directions for adjusting the spreader settings such that the spreader delivers the appropriate application rate recommended above. Apply this product uniformly over the lawn or ornamental turf area. A more uniform application can be made by spreading half of the required amount over the area and then applying the remaining half at a right angle to the previous direction. Avoid streaking, skips, or overlaps during application. Check equipment frequently to ensure equipment is functioning properly and applying uniform distribution of granules.

SUGGESTED SPREADER SETTINGS			
SPREADER	GROUND SPEED	WIDTH OF COVERAGE	
		LOW RATE	HIGH RATE
La. active ingredient/acre (Oxadiazon, Dithiopyr)		1.50 - 0.1875	2.00 - 0.25
Andersons AccoPro 2000, SR 2000 (cone B)	3 mph	12 feet	J 1/2 K 1/2
Andersons AccoPro 1000	3 mph	12 feet	J 1/2 K 1/2
Scotts RSA (cone B)	3 mph	12 feet	H 1/2 I 1/2
Lady Mowls WTR, WFR, HR, (250 PFD at 450 mph) Patelli 11	4.5 mph	36 feet	4 1/2 5 1/4
Vicon (03 Series)	4.5 mph	26 feet	18 20

NOTE: These settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. It is recommended that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic waste supplies.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If items are not acceptable, return the unopened product container at once. To the extent consistent with applicable law, by using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.

WARRANTY DISCLAIMER AND LIMITATION OF LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the Directions. To the extent consistent with applicable law, Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL LIMITED: THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION TO STATE/JURISDICTION. (R19e032105)

Product of USA

Manufactured by:
 The Andersons Lawn Fertilizer Division, Inc.
 P.O. Box 119
 Maumee, OH 43537
 www.andersongolfproducts.com

Andersons Golf Products is a registered trademark of The Andersons, Inc.; Dimension is a registered trademark of Dow AgroSciences LLC; Ronstar is a registered trademark of Bayer.

SGN 125
Net Wt. 50 Lbs. (22.68 kg)

AGCDMR5
 J7 (AGCBKPBRRN.1B)