

Fertilizer with Team* Pro Herbicide 4-4-25



Professional Turf™
Products

FEATURES & USE TIPS:

- Combination granular fertilizer and Team* Pro preemergent herbicide.
- Provides preemergent control of crabgrass and other listed annual grasses and broadleaf weeds in established turfgrasses.
- Consistently sized particles reduce segregation potential.
- Low nitrogen, high potassium formula to build turf tolerance.

PRODUCT SPECIFICATIONS:

N (4%):	1.56% Ammoniacal Nitrogen 2.44% Urea Nitrogen
P - as P₂O₅ (4%):	Ammonium Phosphate
K - as K₂O (25%):	Potassium Chloride
Cl (21.12%):	Maximum chlorine
Active Ingredients:	0.43% Benefin 0.43% Trifluralin

PRODUCT CODE:

APT4T865.2

Date: 080806
Supersedes: 000000
Label: G06

PRODUCT CHARACTERISTICS:

Product Type:

Combination fertilizer &
Team* Pro Herbicide

Nitrogen Type:

Urea

Particle Size:

SGN: 215

Net Weight:

50 lbs.

Coverage:

12,500 sq. ft. @
Low Rate

Bags / Pallet:

45

*Team is a registered trademark of Dow AgroSciences L.L.C.;
The Andersons logo, NS-52 and Professional Turf are trademarks of The Andersons Agriservices, Inc.

*To order, contact your local
Andersons Professional Turf
Products distributor.*

*For more information, contact
Andersons at our website:
www.andersonslawnproducts.com
or call 1-800-225-2639.*

Specimen label follows

PRODUCT SHEET

Fertilizer with Team* Pro Herbicide

4-4-25

A Granular Fertilizer Containing Team Pro Herbicide for the Preemergence Control of Crabgrass, Foxtail, Goosegrass, Poa annua, and Other Annual Grass and Broadleaf Weeds in Established Turfgrass

HERBICIDE

ACTIVE INGREDIENTS:

- Benflin: M-buty-N-ethyl-alpha, alpha, alpha-trifluoro-2,6-dinitro-p-toluidine..... 0.43%
- Trifluralin: alpha, alpha, alpha-trifluoro-2,6-dinitro-N-(4-chlorophenyl)-p-toluidine..... 99.14%
- INERT INGREDIENTS:**..... 100.00%
- Total:**.....

Guaranteed Analysis

- Total Nitrogen (N)..... 4.00%
- 1.58% Ammoniacal Nitrogen
- 2.44% Urea Nitrogen
- Available Phosphate (P₂O₅)..... 4.00%
- Soluble Potash (K₂O)..... 25.00%
- Plant nutrients derived from ammonium phosphate, urea, and potassium chloride.
- Chlorine (Cl) Max..... 21.12%

F292

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994.

EPA Reg. No. 62719-289-9198

EPA Est. No. 9198-OH-28, 9198-AL-001A

Undefined letter is first letter used in run code

*Trademark of Dow AgroSciences LLC

KEEP OUT OF REACH OF CHILDREN WARNING

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes Substantial But Temporary Eye Injury. Harmful If Absorbed Through The Skin.

Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

Personal Protective Equipment

Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical-resistant gloves, shoes plus socks, protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

- Users should:
 - Wash hands after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994, or emergency medical treatment information.

Environmental Hazards: This pesticide is extremely toxic to freshwater marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Read all directions carefully before applying this product.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Entry Restrictions: Do not enter or allow others to enter the treated area until dusts have settled. If watering-in is required after application, do not enter or allow others to enter the treated area (except those involved in the watering-in) until the watering-in is complete and the surface is dry.

Storage and Disposal: Do not contaminate water, food, or feed by storage or disposal.
Pesticide Storage: Store in original container only. In case of a leak or spill, contain material and dispose as waste.
Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

General Information
 This fertilizer product contains Team Pro, a preemergence herbicide for control of annual grasses and broadleaf weeds in established turfgrasses.

Annual Grasses:	Scientific Name	Common Name
crabgrass, large	<i>Digitaria sanguinalis</i>	Spencer crabgrass
crabgrass, smooth	<i>Digitaria sanguinalis</i>	Spencer crabgrass
crabgrass, large or hairy	<i>Digitaria sanguinalis</i>	Spencer crabgrass
goosegrass (silver crabgrass or cowbirdgrass)	<i>Elytrigia inodora</i>	Eleusine indica
green foxtail	<i>Echinochloa crus-galli</i>	Echinochloa crus-galli
green foxtail	<i>Echinochloa crus-galli</i>	Echinochloa crus-galli
yellow foxtail	<i>Setaria viridis</i>	Setaria viridis
Broadleaf Weeds:	Scientific Name	Common Name
chickweed, common	<i>Stellaria media</i>	Stellaria media
knotted gramine	<i>Polygonum aviculare</i>	Polygonum aviculare
lambquarters, common	<i>Chenopodium album</i>	Amaranthus spp.
pigweed	<i>Amaranthus spp.</i>	Amaranthus spp.
purslane, common	<i>Portulaca oleracea</i>	Portulaca oleracea
Weeds Suppressed:	Scientific Name	Common Name
Common bermuda	<i>Cynodon dactylon</i>	Common bermuda
Florida betelnut	<i>Richardia scabra</i>	Richardia scabra
prostrate sungrape	<i>Euphorbia humifusa</i>	Euphorbia humifusa
spotted sungrape	<i>Euphorbia maculata</i>	Euphorbia maculata
creeping sungrape	<i>Euphorbia serpens</i>	Euphorbia serpens
creeping woodsorel	<i>Oxalis corniculata</i>	Oxalis spp.
woodsorel	<i>Oxalis spp.</i>	Oxalis spp.

Uses

Established Turfgrass
When to Apply: For optimum results, this product must be applied one to two weeks prior to the anticipated germination of the target annual weeds. Application will usually provide season-long weed control. However, a second or split application is recommended when the initial application is made well in advance of weed germination or when an extended period of weed control is required, such as areas with longer growing seasons or for lawn care operator programs.

How to Apply: Apply this product evenly over the turfgrass area using a properly calibrated drop type or rotary spreader. Calibrate the spreader prior to use according to the manufacturer's directions. Check frequently to make sure equipment is working properly and distributing granules uniformly. More uniform application may be achieved by spreading half of the required amount of product over the area and then applying the remaining half in swaths at a right angle to the first. Avoid skips and excessive overlaps as poor weed control or turfgrass injury may occur. Avoid use of spreaders that apply material in narrow concentrated bands. A splash board mounted below the release point of drop-type spreaders can improve the uniformity of product distribution.

Activation: Optimum weed control performance will be obtained if treated areas receive rainfall or irrigation soon after application.

Important Use Precautions: Do not apply this product to golf course putting greens, dionchroa, or newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass. Any cultural practices that disturb the soil, such as aeration or verticiling, should be done prior to application. This product will not harm most nearby ornamentals.

This product may be applied to the following established turfgrass species:

Established Cool Season Turfgrasses	Established Warm Season Turfgrasses
Bentgrass fairways	Bahagrass
Kentucky bluegrass	Bermudagrass
Perennial ryegrass	Centipedegrass
Fescues	Fescue, tall
	St. Augustinegrass
	Zoysiagrass

Broadcast Application Rates:	Application Rate	Specific Use Directions
Application Timing: For extended control of spring and summer annual weeds.	Spring: 174 to 349 lbs/acre (4.00 to 8.01 lbs/1,000 sq. ft.) Fall: 174 lbs/acre followed by 4.00 lbs/1,000 sq. ft.	In cool season and warm season turfgrasses, make a single application in the spring or fall, 1 to 2 weeks before expected time of weed germination. In cool season turfgrass area and northern half of the warm season turfgrass area, apply the first application in spring 1 to 2 weeks before expected application and a second application 8 to 10 weeks later.
Split Application: For extended control of spring and summer annual weeds.	233 lbs/acre followed by 233 lbs/acre (5.34 followed by 5.34 lbs/1,000 sq. ft.)	In southern half of warm season turfgrass area, apply the first application in spring 1 to 2 weeks before the expected time of weed germination and make a second application 10 to 12 weeks later.

Reseeding Precautions
 Team Pro herbicide, which controls annual grass weeds, may also affect the establishment of new desirable turfgrass seedlings, if reseeding is attempted too soon after application. When reseeding, it is essential that proper cultural practices, such as soil cultivation, irrigation and fertilization be followed. Reseeding, following the last application of this product, should not be attempted until the applicable minimum time delay in the following table has been met:

Amount of product Applied Per Acre	Minimum Time Delay Following Last Application Before Reseeding (Weeks)
Single application of 174 lbs/acre (4.00 lbs/1,000 sq. ft.)	8
Single application of 349 lbs/acre (8.01 lbs/1,000 sq. ft.)	12 to 16
Split application of 174 lbs/acre (4.00 lbs/1,000 sq. ft.)	8 to 10
Split application of 233 lbs/acre (5.34 lbs/1,000 sq. ft.)	10 to 12

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			LOW RATE	MEDIUM RATE	HIGH RATE
Lb. active ingredient/acre (benflin, trifluralin)			0.75, 0.75	1.00, 1.00	1.50, 1.50
Andersons AccuPro 2000, SR-2000 (cone 3)	3 mph	12 feet	K 12	L 12	N 14
Andersons AccuPro 1000	3 mph	12 feet	K 12	L 12	N 14
Scotts R-8A (cone 3)	3 mph	11 feet	L 14	M	O
Lely Models WTR, WFR, HR, 1250 (P 10 at 450 rpm)	4.5 mph	52 feet	5 14	6 14	8 14
Vicon (03 series)	4.5 mph	38 feet	26	31	42

Terms and Conditions of Use
 If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

Warranty Disclaimer
 Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use
 It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as intolerable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Seller. All such risks shall be assumed by Buyer.

Limitation of Remedies
 The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal agreements or agreements. No employee or sales agent of the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Distributed by:
 The Andersons Lawn Fertilizer Division, Inc.
 PO Box 119
 Maumee, OH 43837

SGN 215

**Net Wt. 50 Lbs. (22.68 kg)
 Covers Up To 12,500 Sq. Ft.**

Contains 0.43 pounds active ingredient per 50 pound bag

G6 (APTBLKPDNAB)

APT4T865.2