



Professional Turf™ Products

1% Bayleton® Fungicide

FEATURES & USE TIPS:

- For control of Dollar Spot, Anthracnose, Brown Patch, Summer Patch, and other diseases as listed on the label for turf.
- Contains Bayleton® branded fungicide.
- Bayleton® fungicide is absorbed rapidly and works systemically from within the plant.

PRODUCT SPECIFICATIONS:

Active Ingredient: Triadimefon (1.0%)

PRODUCT CODE:

APTBY11.1

Date: 071007
Supersedes: 000000

PRODUCT CHARACTERISTICS:

Product Type:
Granular Fungicide

Particle Size:
SGN: 125

Net Weight:
15 lbs.

Coverage:
10,000 sq. ft @
Low Rate

Bags / Pallet:
96

The Andersons logo is a registered trademark of The Andersons, Inc. Professional Turf is a trademark of The Andersons, Inc. Bayleton is a registered trademark of Bayer.

To order, contact your local Andersons Golf Products distributor.

*For more information, contact Andersons at our website:
www.andersonsgolfproducts.com
or call 1-800-225-2639.*

Following is a specimen label that may at some point be revised or otherwise become out of date. It is intended as a guide in providing general information regarding the use of this product. Always read and follow the label on the product container.



Professional Turf™ Products

1% Bayleton® Fungicide

FOR CONTROL OF CERTAIN DISEASES ON TURF

ACTIVE INGREDIENT:

Triadimenol* 1.0%

OTHER INGREDIENTS 99.0%

Total 100.0%

*CAS No. 43121-43-3

Bayleton is a registered trademark of Bayer

EPA Reg. No. 9198-111

EPA Est. No. 9198-OH-1M, 9198-OH-2B, 9198-AL-001A

Underlined letter is first letter used in run code on bag.

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing vapor or dust. Avoid contact with skin, eyes or clothing. Do not contaminate feed or food. Keep out of reach of children.

Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before reuse.

If swallowed

Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in eyes

Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contacts, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If inhaled

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-757-8851 for emergency medical treatment information.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as seed, or for commercial seed production, or for research purposes.

APPLICATION: Bayleton fungicide is absorbed rapidly and works systemically from within the plant. Rainfall or sprinkler irrigation soon after application will assure rapid systemic uptake. Control will be less effective on turf suffering from drought stress. Therefore, in order to achieve maximum control, plants should be maintained in a vigorously growing state through good cultural practices.

Apply when the turf is fully established and actively growing. Application should be applied at recommended intervals to maintain disease control.

RECOMMENDED APPLICATIONS TURF GRASS DISEASE CONTROL:

DISEASE	Lbs. of Product Per 1,000 sq. ft.*		REMARKS
	PREVENTATIVE	CURATIVE	
Dollar Spot	1.5	--	PREVENTATIVE RATE: Apply recommended rate on 30-day intervals. Protective activity of Bayleton fungicide may extend for as long as 60 days dependent upon environmental conditions.
	0.75	--	PREVENTATIVE RATE: Apply recommended rate on 14-day intervals. Protective activity of Bayleton fungicide may extend for as long as 30 days dependent upon environmental conditions.
	--	3.0	CURATIVE RATE: To control existing infections, apply the curative rate. Re-application should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedules for rates and timing.
Copper Spot Powdery Mildew Red Thread (<i>Laetisaria fulcriformis</i>) Risks Brown Patch (<i>Rhizoctonia solani</i>) (Suppression)	1.5	--	PREVENTATIVE RATE: Apply at 15 to 30 day intervals. When environmental conditions favor light-to-moderate disease development, use a longer interval. Protective activity of Bayleton fungicide can be greater than 30 days depending on environmental conditions.
	--	3.0	CURATIVE RATE: To control existing infections, apply the curative rate. Re-applications should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedule for rates and timing.
Antracnose <i>Poa annua</i> /decline	3.0	--	PREVENTATIVE RATE: Apply at 30-day intervals and repeat as necessary for seasonal control. Dependent upon environmental conditions, residual control may be extended to 45 days.
	1.5	--	PREVENTATIVE RATE: Apply at 21-day intervals and repeat as necessary for seasonal conditions, residual control may be extended to 30 days.
	--	3.0	CURATIVE RATE: To control existing infections, apply the curative rate. Re-applications should be no closer than 14 days. When infections have been controlled, refer to the preventative rate schedules for rates and timing.
	PREVENTATIVE RATES ONLY		
Stripe Smolt	3.0		Make a single application in spring as grass growth begins.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>) Summer Patch (<i>Phialophora graminicola</i>) (Except California)	3.0		Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at 30-day intervals.
Fusarium Blight (<i>Fusarium culmorum</i>) (<i>Fusarium poae</i>) Summer Patch (<i>Phialophora graminicola</i>) (California only)	3.0 to 6.0		Apply first application in mid-June or 30 days prior to time this blight normally becomes evident. Applications should be made at the 6 b. rate on 15-day intervals.
Gray Snow Mold (<i>Typhula incarnata</i>) (Except California)	6.0		Apply in fall 30 days prior to turf dormancy. If turf breaks dormancy during winter months, a second application should be made. Do not apply over snow cover or when turf is dormant.
Pink Snow Mold (<i>Microbotryum male</i>) Southern Blight (<i>Sclerotium oryzae</i>)	3.0 to 6.0		Apply before conditions favorable for infection occur. Reapplication should be made as needed at 60 to 90-day interval. Do not apply over snow cover or when turf is dormant. Use higher rate in areas with a history of severe disease damage.
Gray Snow Mold (<i>Typhula incarnata</i>) Pink Snow Mold (<i>Microbotryum male</i>) (California Only):	Single Application 7.5 to 12.0		Apply in late fall before first snow cover. Do not apply over snow cover. Use the high rate in areas with a history of severe snow mold damage.

NOTE*: Apply the specified amount of this product with a standard granular applicator accurately calibrated to uniformly apply recommended amount of product per 1,000 sq. ft.

Make application when turf is dry and following the mowing operation. Do not mow until a minimum of 0.25 inch of water has been applied to treated area. Do not use clippings for animal feed.

SUGGESTED SPREADER SETTINGS

These suggested spreader settings are not intended to replace calibration. Please calibrate your spreader before applying product.

15 lbs. treats 10,000 sq. ft. (0.23 acre) at the 65 lbs. product/acre - LOW RATE (1.5 lbs/1,000 sq. ft.)
15 lbs. treats 5,000 sq. ft. (0.11 acre) at the 130 lbs. product/acre - HIGH RATE (3.0 lbs/1,000 sq. ft.)

SPREADER	Ground Speed	Width of Coverage	LOW RATE 0.65 lbs. ai/acre	HIGH RATE 1.30 lbs. ai/acre
Andersons AccuPro 2000, SP-2000 (cone 6)	3 mph	6 feet	1	1 1/2
Andersons AccuPro 1000	3 mph	6 feet	1	1 1/2
Andersons SS-2 Drop	3 mph	overlap wheels	3 1/4	4
Scotts R-84 (cone-6)	3 mph	6 feet	H	H 1/2
Lely Models WTR, WFR, HR, 1250 (PTO at 450 rpm), Pattern I/C	4.5 mph	15 feet	3 1/2	4
Vicon (09 Series)	4.5 mph	10 feet	16	20

NOTE: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area to be treated. Measure area to be treated. Do not over or under apply. Close spreader when filling, turning or stopping. DO NOT HAND SPREAD.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feed-stuffs, and domestic water supplies.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State or local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or Buyer accepts the following Warranty Disclaimer and Limitation of Liability.

WARRANTY DISCLAIMER and LIMITATION of LIABILITY:

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. Manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Where permissible, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS. YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION. (AP21993/RY/rev060304)

Attention: This product contains a chemical known to the State of California to cause birth defects or other reproductive harm.

Net Wt. 15 Lbs. (6.8 kg)
Treats Up To 10,000 Sq. Ft.

Manufactured by:
The Andersons Lawn Fertilizer Division, Inc.
P.O. Box 119, Maumee, OH 43537
www.andersonslawnproducts.com

G6 (BLKP1326B)

APTBY11.1